



WATER STRUCTURE AND SCIENCE


Martin Chaplin



Anomalous properties of water

Many of the properties of water are quite different from those expected from those of other liquids.

- ▼ [The range of anomalous properties of water](#)
- ▼ [Rationale for the low-temperature anomalies of liquid water](#)
- ▼ [Water phase anomalies](#)
- ▼ [Water density anomalies](#)
- ▼ [Water material anomalies](#)
- ▼ [Water thermodynamic anomalies](#)
- ▼ [Water physical anomalies](#)
- ▼ [Pure water](#)



'...water has the remarkable property of being more compressible in winter than in summer..'

John Canton 1764^j

The range of anomalous properties of water

The anomalous properties of water are those where the behavior of liquid water is quite different from what is found with other liquids [1414].^a The strength and directionality of the **hydrogen bond** are control liquid water's thermodynamic and dynamic behavior. If hydrogen-bonding did not exist, water would behave non-anomalous as expected from similar molecules. No other material is commonly found as solid (ice), liquid (water), or gas (steam).^d Frozen water (ice) also shows anomalies when compared with other solids. Although it is an apparently simple molecule (H₂O), it has a highly complex and anomalous character due to its inter-molecular **hydrogen bonding** (see [1530] for example). As a gas, water is one of **lightest known**, as a liquid, it is much **denser than expected** and as a solid, it is much **lighter than expected** when compared with its liquid form. It can be extremely slippery and extremely sticky at the same time^d, and this 'stick/slip' behavior is how we recognize the feel of water [2411]. Many other anomalies of water may remain to be discovered, such as the possible link of water to room temperature superconductivity [2124].^h An interesting history of the study of the anomalies of water has been published [1542].

As liquid water is so common-place in our everyday lives, it is often regarded as a 'typical' liquid. In reality, water is most atypical as a liquid, behaving as a quite different material at low temperatures to that when it is hot, with a division temperature of about 50 °C. It has often been stated (for example, [127]) that life depends on these anomalous properties of water. The anomalous macroscopic properties of water are derived from its microscopic structuring.

The high cohesion between molecules gives it a high freezing and melting point, such that we and our planet are bathed in liquid water. The large heat capacity, high thermal conductivity and high water content in organisms contribute to thermal regulation and prevent local temperature fluctuations, thus allowing us to more easily control our body temperature. The high latent heat of evaporation gives resistance to dehydration and considerable evaporative cooling. It has unique **hydration properties** towards important biological macromolecules (particularly proteins and nucleic acids) that determine their three-dimensional structures, and hence their biological functions, in solution. This hydration forms gels that can reversibly undergo the gel-sol phase transitions that underlie many cellular mechanisms [351]. **Water ionizes** and allows easy proton exchange between molecules, so contributing to the richness of the ionic interactions in biology. Also, it is an **excellent solvent** due to its polarity, high relative permittivity (dielectric constant) and small size, particularly for polar and ionic compounds and salts.

At 4 °C water expands on **heating or cooling**. This density maximum together with the low ice density results in (i) the necessity that all of a body of fresh water (not just its surface) is close to 4 °C before any freezing can occur, (ii) the freezing of rivers, lakes, and oceans is from the top down, so permitting survival of the bottom ecology, insulating the water from further freezing, reflecting back sunlight into space and allowing rapid thawing, and (iii) density driven thermal convection causing seasonal mixing in deeper temperate waters carrying life-providing oxygen into the depths. The large heat capacity of the oceans and seas allows them to act as heat reservoirs such that sea temperatures vary only a third as much as land temperatures and so moderate our planet's climate (for example, the Gulf stream carries tropical warmth to northwestern Europe). The compressibility of water reduces the sea level by about 40 m giving us 5% more land [65]. Water's high **surface tension** plus its **expansion on freezing** encourages the erosion of rocks to give soil for our agriculture.

Notable amongst the anomalies of water is the opposite properties of hot and cold water, ^b with the anomalous behavior more accentuated at low temperatures where the properties of supercooled water often diverge from those of hexagonal ice. In particular, several properties of water change at about 50 °C [2755]; just above the body temperature of mammals and about which many proteins denature. As (supercooled) cold liquid water is heated **individual molecules shrink**, bulk water **shrinks** and becomes less easy to **compress**, its **refractive index** increases, the **speed of sound** within it increases, gases become **less soluble** and it is **easier to heat** and **conducts heat** better. In contrast, as hot liquid water is heated it **expands**, it becomes easier to **compress**, its **refractive index** reduces, the **speed of sound** within it decreases, gases become **more soluble** and it is **harder to heat** and a poorer **conductor of heat**. With increasing pressure, **individual molecules expand**, cold water molecules **move faster** but hot water molecules **move slower**. Hot water **freezes faster** than cold water and ice **melts when compressed** except at high pressures when liquid water **freezes when compressed**.

Fluctuations in liquid water ⁱ

$$\begin{aligned}\langle(\Delta V)^2\rangle_{\text{TPN}} &= T \times \kappa_T \times V \times k_B \\ \langle(\Delta V)(\Delta S)\rangle_{\text{TPN}} &= T \times \alpha_P \times V \times k_B \\ \langle(\Delta S)^2\rangle_{\text{TPN}} &= N \times C_P \times k_B\end{aligned}$$


increase upon cooling below a certain temperature, and down to the homogeneous ice nucleation temperature (~ -45 °C), instead of decreasing as expected.

The anomalies of water appear as a hierarchy of effects with different bounds [169]. These are shown indicatively opposite as derived from modeling, not experimental data. The 'Structural' bounds indicate where water is more disordered when compressed, the 'Dynamic' bounds indicate where **diffusion increases with density**, and the 'Thermodynamic' bounds show where there is a **temperature of maximum density**.⁹ As density always increases with increasing pressure, a similar relationship holds with pressure along the horizontal axis.

Phase diagrams of other tetrahedrally structured liquids (e.g. Si, SiO₂) also show the nesting of anomalous regions [2285].

Other anomalies of water are caused by the onset of interpenetration at about 200 MPa, such as the **high-pressure C_p anomaly** [2929], the **density-distance paradox** and the **fast-sound anomaly**.

[[Back to Top](#) ▲]



Rationale for the low-temperature anomalies of liquid water

Four different scenarios have been suggested:

(1a) **The stability limit conjecture.** There is a limit (a singularity) to the mechanical stability of water at -45°C [1886, 2995]. This was suggested to be associated with the cooperative formation of an open hydrogen-bonded network.

(1b) **A singularity-free scenario.** It was suggested that the simplest interpretation of the behavior of supercooled water consistent with experimental observations is free of singularities [2996] This is consistent with supercooled water approaching the structure of a fully bonded random or non-random (e.g. **ES**) tetrahedral network on cooling. This may or may not be linked to a critical point at higher or negative pressures.

(2a) **A liquid-liquid critical point.** Increasingly, scientists attribute the low-temperature anomalous nature of water to the presence of a metastable **second critical point** at about -50°C (under pressure) when high-density liquid water and low-density liquid water no longer coexist within the same phase [2930, 3134] but split into different phases, separated by a first-order phase transition.

(2b) **A critical point free scenario.** The "fragile-to-strong" transition for supercooled water, rather than a critical point, is interpreted as the reason for the anomalies in supercooled water [3121].

It has recently been established that a **second critical point** exists [2930, 3134] and the data supporting the other scenarios are all consistent with 2a above. Water's anomalies do not require this as an explanation, but it does seem likely [2947] that there is such a phenomenon and it would cause the attributed effects. The liquid-liquid critical point scenario does not contain any information concerning the structure of the two phases involved and, in this respect, it is a

somewhat unproductive hypothesis as a sole explanation of these anomalies (as the attribution mixes cause with effect, as agreed by others [1859]). Description of the structures is [given elsewhere](#).

[\[Back to Top ▲\]](#)

Water phase anomalies ^d

1. Water has an unusually high [melting point](#). [[Explanation](#)]
2. Water has an unusually high [boiling point](#). [[Explanation](#)]
3. Water has an unusually high [critical point](#). [[Explanation](#)]
4. [Solid water](#) exists in a wider variety of stable (and metastable) crystal and amorphous structures than other materials. [[Explanation](#)]
5. The thermal conductivity, shear modulus and transverse sound velocity of ice reduce with increasing pressure. [[Explanation](#)]
6. The structure of liquid water changes at high pressure. [[Explanation](#)]
7. Supercooled water has two phases and a [second critical point](#) at about -91 °C. [[Explanation](#)]
8. Liquid water is easily supercooled but glassified with difficulty. [[Explanation](#)]
9. Liquid water exists at very low temperatures and freezes on heating. [[Explanation](#)]
10. Liquid water may be easily superheated. [[Explanation](#)]
11. Hot water may freeze faster than cold water; the Mpemba effect. [[Explanation](#)]
12. Warm water vibrates longer than cold water. [[Explanation](#)]
13. Water molecules shrink as the temperature rises and expand as the pressure increases. [[Explanation](#)]

[\[Back to Top ▲\]](#)

Water density anomalies

1. The density of ice increases on heating (up to 70 K). [[Explanation](#)]
2. Water shrinks on melting. [[Explanation](#)]
3. Pressure reduces ice's melting point. [[Explanation](#)]
4. Liquid water has a high-density that increases on heating (up to 3.984 °C). [[Explanation](#)]
5. The surface of water is denser than the bulk. [[Explanation](#)]
6. Pressure reduces the temperature of maximum density. [[Explanation](#)]
7. There is a minimum in the density of supercooled water. [[Explanation](#)]
8. Water has a low coefficient of expansion (thermal expansivity). [[Explanation](#)]
9. Water's thermal expansivity reduces increasingly (becoming negative) at low temperatures. [[Explanation](#)]
10. Water's thermal expansivity increases with increased pressure. [[Explanation](#)]
11. The number of nearest neighbors increases on melting. [[Explanation](#)]
12. The number of nearest neighbors increases with temperature. [[Explanation](#)]
13. Water has unusually low compressibility. [[Explanation](#)]
14. The compressibility drops as temperature increases up to 46.5 °C. [[Explanation](#)]
15. There is a maximum in the compressibility-temperature relationship. [[Explanation](#)]
16. The speed of sound increases with temperature up to 74 °C. [[Explanation](#)]
17. The speed of sound may show a minimum. [[Explanation](#)]
18. 'Fast sound' is found at high frequencies and shows a discontinuity at higher pressure. [[Explanation](#)]
19. NMR spin-lattice relaxation time is very small at low temperatures. [[Explanation](#)]
20. The NMR shift increases to a maximum at low (supercooled) temperatures [[Explanation](#)]
21. The refractive index of water has a maximum value at just below 0 °C. [[Explanation](#)]
22. The change in volume as liquid changes to gas is very large. [[Explanation](#)]

[\[Back to Top ▲\]](#)

Water material anomalies

1. No aqueous solution is ideal. [[Explanation](#)]
2. D₂O and T₂O differ significantly from H₂O in their physical properties. [[Explanation](#)]
3. Liquid H₂O and D₂O differ significantly in their phase behavior. [[Explanation](#)]
4. H₂O and D₂O ices differ significantly in their quantum behavior. [[Explanation](#)]
5. The mean kinetic energy of water's hydrogen atoms increases at low temperature (disputed). [[Explanation](#)]
6. Solutes have varying effects on properties such as density and viscosity. [[Explanation](#)]
7. The solubilities of non-polar gases in water decrease with temperature to a minimum and then rise. [[Explanation](#)]
8. The dielectric constant of water and ice are high. [[Explanation](#)]
9. The relative permittivity shows a temperature maximum. [[Explanation](#)]
10. The relative permittivity shows a 'kink' in its behavior with the temperature at 60 °C. [[Explanation](#)]
11. The imaginary part of the dielectric constant shows a minimum near 20 K. [[Explanation](#)]
12. Proton and hydroxide ion mobilities are anomalously fast in an electric field. [[Explanation](#)]
13. The electrical conductivity of water rises to a maximum at about 230 °C. [[Explanation](#)]

14. The electrical conductivity of water rises considerably with frequency. [Explanation]
15. Acidity constants of weak acids show temperature minima. [Explanation]
16. X-ray diffraction shows an unusually detailed structure. [Explanation]
17. Under high pressure water molecules move further away from each other with increasing pressure; a density-distance paradox. [Explanation]
18. Water adsorption may cause negative electrical resistance. [Explanation]

[Back to Top ▲]

Water thermodynamic anomalies


1. The heat of fusion of water with temperature exhibits a maximum at -17 °C. [Explanation]
2. Water has over twice the **specific heat** capacity of ice or steam. [Explanation]
3. The **specific heat** capacity (C_p and C_v) is unusually high. [Explanation]
4. The specific heat capacity C_p has a minimum at 36 °C. [Explanation]
5. The **specific heat** capacity (C_p) has a maximum at about -45 °C. [Explanation]
6. The **specific heat** capacity (C_p) has a minimum with respect to pressure. [Explanation]
7. The **heat** capacity (C_v) has a maximum. [Explanation]
8. The high **heat of vaporization**. [Explanation]
9. The high heat of sublimation. [Explanation]
10. The high entropy of vaporization. [Explanation]
11. The thermal conductivity of water is high and rises to a maximum at about 130 °C. [Explanation]

[Back to Top ▲]

Water physical anomalies

1. Water has unusually high **viscosity**. [Explanation]
2. Large **viscosity** and Prandtl number increase as the temperature is lowered. [Explanation]
3. Water's viscosity decreases with pressure below 33 °C. [Explanation]
4. Large diffusion decrease as the temperature is lowered. [Explanation]
5. At low temperatures, the self-diffusion of water increases as the density and pressure increase. [Explanation]
6. The thermal diffusivity rises to a maximum at about 0.8 GPa. [Explanation]
7. Water has unusually high **surface tension**. [Explanation]
8. Some salts give a surface tension-concentration minimum; the Jones-Ray effect. [Explanation]
9. Some salts prevent the coalescence of small bubbles. [Explanation]
10. The molar ionic volumes of salts show maxima with respect to temperature. [Explanation]

[Back to Top ▲]



The figure left shows some of the anomalous properties of liquid water that are related to temperature. The graph uses data that have been scaled between their maximum and minimum values within this range (see [original data](#)). Note, in particular, the different behaviors at high and low temperatures that is in marked contrast to the behavior of simple liquids, whose thermodynamic properties vary in a monotonic manner with temperature.

Footnotes

^a Whether or not the properties of water are seen to be anomalous depends upon the materials used in the comparison and the interpretation of the term 'anomalous'. For example, it could well be argued that water possesses exactly those properties that one might deduce from its structure (see for example, [402]). The small number of other tetrahedrally

interacting liquids (without any hydrogen bonds), such as liquid Si [1835], SiO₂ and BeF₂ have many similar 'anomalies' [1814], as do other materials (also without any hydrogen bonds) where mixed phases may arise, such as liquid Te [1538]. Comparisons between water, liquid sodium, argon, and benzene appeared to Franks [112] to indicate several of the properties given above as not being anomalous. However, these materials are perhaps not the most typical of liquids and elsewhere Franks endorses water's anomalous nature. Also, other liquids (e.g. hydrazine, N₂H₄, melting point 2 °C, boiling point 114 °C) are multiply hydrogen-bonded but do not show similar anomalies. My list gives the unusual properties generally understood to make liquid water (and ice) stand out from 'typical' liquids (or solids). See [242] for a review concentrating on the non-anomalous properties of water; that is, those that are the 'same' as for other liquids. At higher temperatures (>315 K) the thermodynamic properties of water may be considered close to 'normal' for a liquid [1638]. Note that properties that are compared at ambient pressure or along the vapor/liquid line may be seen as anomalous whereas under isopycnic (isodensity) conditions no (anomalous) maximum or minimum values may be found (for example, no **specific heat minimum**, **speed of sound maximum** or **compressibility minimum** are found at constant water density of 1 g cm⁻³; see [line on phase diagram](#)).


It is generally agreed by water scientists today that water is the most anomalous substance [2653]. [\[Back\]](#)

b Some scientists attribute the low-temperature anomalous nature of water to the presence of a **second critical point**; an interesting if somewhat unproductive hypothesis as a sole explanation (as the attribution mixes cause with effect, as agreed by others [1859]). Water's anomalies do not require this as an explanation, although it does seem likely [2947]. [\[Back\]](#)

c The temperature range of 'hot' and 'cold' water varies in these examples; see the individual entries for details. [\[Back\]](#)

d The anomalies of water are divided into groups but, clearly, some anomalies may be included under more than one topic and there may not be a universal agreement for the groupings shown. The 'number' of anomalies depends on which ones are chosen and whether related anomalies (such as reciprocal variations including molar volume and density) are grouped together or as separate phenomena. [\[Back\]](#)

e This is easily shown with two wet glass panes. If one sheet is placed horizontally on top of the other then the sheets easily slip over each other horizontally (i.e. very slippery) whilst it being almost impossible to separate the sheets vertically (i.e. very sticky). A further example of its stickiness is its use in making sand castles, whereas its slipperiness is well known on wet floors. The stickiness of water ice in the outer reaches of protoplanetary disks enhances the growth of water-containing rocky asteroids and planets as the particles stick together [2251]. [\[Back\]](#)



f The data from [169] is shifted upwards 38 K to give the correct temperature of maximum density under standard pressure. [\[Back\]](#)

g Sometimes apparently unpredictable or unexpected properties of liquid water may be due to variations in the dissolved gas concentrations [1948], a factor that is difficult to control and easy to overlook. Atmospheric gases dissolve in water and then form nanobubbles and microbubbles some of which may expand and rise back to the surface. This process causes continuous but somewhat chaotic, changes in the gaseous concentrations over significant time periods (> 100 s) and consequently continuous changes in the hydrogen-bonded structuring within the water [1948]. Such artifacts are thought to be absent in the anomalous properties described above. [\[Back\]](#)

h Where $\langle(\Delta V)^2\rangle_{TPN}$ is the mean square fluctuations in volume (V) for constant temperature (T), pressure (P) and number of molecules (N), $\langle(\Delta V)(\Delta S)\rangle_{TPN}$ is the fluctuation correlation in volume and entropy for constant temperature, pressure and number of molecules, $\langle(\Delta S)^2\rangle_{TPN}$ is the mean square fluctuations in entropy for constant temperature, pressure and number of molecules, κ_T is the isothermal compressibility, k_B is the **Boltzmann constant**, α_P is the thermal expansivity and C_P is the heat capacities at constant pressure. [\[Back\]](#)

i J. Canton, Experiments and observations on the compressibility of water and some other fluids, *Phil. Trans.* **54** (1764) 261-262. [\[Back\]](#)

[Home](#) | [Site Index](#) | [Explanation of the anomalies](#) | [Water: Introduction](#) | [The icosahedral water clusters](#) | [LSBU](#) | [Top](#)

This page was established in 2003 and last updated by [Martin Chaplin](#) on 4 January, 2018



This work is licensed under a [Creative Commons Attribution-NonCommercial-No Derivative Works 2.0 UK: England & Wales License](#)